



Dr. Mikkel C. Jarman, DPM

633 E. Ray Rd. Suite 128  
Gilbert, AZ 85296

ph: 480-497-3946

fx: 480-497-3947

## WART SURGERY SAME DAY

### INSTRUCTIONS FOR WOUND CARE:

1. Numbness will last for 1 (one) to 4 (four) hours, possibly longer.
2. White streaking in the foot will quickly disappear as the anesthetic is absorbed.
3. Some bleeding and oozing is normal, but if excessive, apply direct constant pressure with clean gauze for fifteen (15) minutes. An ice pack is also helpful. If bleeding is persistent, call the office.
4. Extra Strength Tylenol or Advil is usually sufficient for discomfort.
5. Limit activities today.

### FOLLOWING MORNING:

1. Remove the dressing. Hydrogen peroxide can be used if difficult to remove. You may then shower.
2. Wash the wound well with soap and water. Do not scrub the site.
3. After the skin is dry, coat the area well with Neosporin Cream, a topical antibiotic that can be purchased without prescription.
4. Cover with a cloth band-aid. Keep it clean and dry.
5. Repeat this process one (1) or two (2) times a day.
6. Limit activities to what you can tolerate. Swimming in a pool is generally allowed, but avoid lakes and rivers.
7. Once the healing process is well underway, the bandage can be removed at night, but keep it covered during the day.
8. Follow-up appointments are very important to monitor healing and assure that no new warts have developed. It usually takes approximately one (1) month or longer for complete healing.
9. After you are discharged, continue to check your feet frequently. Because warts are caused by a virus, new warts can appear and are the easiest to remove when small. If left untreated, warts can spread.
10. Avoid going barefoot in public places such as showers and locker rooms. The wart virus spreads easily in moist conditions, so wear thongs. Protect your feet. Keep them clean and dry.

**For additional advice or concerns, please contact our office, 480-497-3946**